

**SOURCE SCALING OF AFTERSHOCKS FOLLOWING THE 2009 MW 6.3
L'AQUILA EARTHQUAKE (CENTRAL ITALY) RECORDED AT AN
UNDERGROUND ARRAY**

Danilo Galluzzo

(INGV- Sezione di Napoli, Osservatorio Vesuviano)

Most of the studies about source parameters of large-to-small earthquakes uses far-field surface recordings from regional seismic network. Seismic noise, site and path effects can strongly bias the earthquake source spectral parameters, particularly for earthquakes of low-to-moderate magnitude. The use of seismic signals recorded in underground environment, where seismic noise is reduced, and the application of Spectral Ratio Approach (SRA) between similar events are strongly recommended for the investigation of source parameters such as stress drop and corner frequency. We show the results obtained on 7500 earthquakes of low-to-moderate magnitude ($0.5 < M_L < 4.1$), recorded at the underground array Underseis located at 1.4 km depth near the town of L'Aquila. First we identified 290 clusters of similar seismic events by applying a cross-correlation based criterion, then we apply the spectral ratios approach among the events of each cluster to estimate the source parameters. The most of clusters are located close to the main shock of April 6, 2009, at a distance of 15-25 km from the array. For each cluster we apply SRA to evaluate source spectral parameters such as the corner frequencies by using a grid search algorithm and consequently the stress drop. The S-wave spectra averaged on 7 stations of the array were used to evaluate spectral ratios. The redundancy of solutions obtained by considering the spectral ratios among all possible pairs of similar events is used for a more stable estimate of the corner frequencies. The seismic moment was estimated from the low frequency part of the averaged S-wave spectra. The stress drop vs seismic moment trend shows non self similar behavior for the investigated range of magnitude ($0.5 < M_L < 4.1$). The apparent stresses clearly increase from 1 to 200 bars with seismic moment increasing from 10^{10} to 10^{14} Nm. Preliminary results show that SRA and array underground data are particularly suited to investigate the spectral source parameters of small earthquakes.